arXiv:astro-ph/9803033v1 4 Mar 1998

Properties of Very Luminous Galaxies

A. Cappi

Osservatorio Astronomico di Bologna, via Zamboni 33, I-40126, Bologna, Italy; cappi@astbo3.bo.astro.it

L.N. da Costa[1]

European Southern Observatory, Karl-Schwarzschild-Str.2, D-85748 Garching bei München, Germany; ldacosta@eso.org

C.Benoist

European Southern Observatory, Karl-Schwarzschild-Str.2, D-85748 Garching bei München, Germany; cbenoist@eso.org

S. Maurogordato

CNRS; Observatoire de Nice, B4229, Le Mont–Gros, 06304 Nice Cedex 4, France; maurogor@faure.obs-nice.fr

P. S. Pellegrini

Departamento de Astronomia CNPq/Observatório Nacional, rua General José Cristino 77, Rio de Janeiro, R.J. 20921 Brazil; pssp@dans.on.br

Received _____; accepted _____

The Astronomical Journal, in press

[1]Departamento de Astronomia CNPq/Observatório Nacional, rua General José Cristino 77, Rio de Janeiro, R.J. 20921 Brazil

ABSTRACT

Recent analysis of the SSRS2 data based on cell–counts and two–point correlation function has shown that very luminous galaxies are much more strongly clustered than fainter galaxies. In fact, the amplitude of the correlation function of very luminous galaxies ($L > L^*$) asymptotically approaches that of $R \geq 0$ clusters. In this paper we investigate the properties of the most luminous galaxies, with blue absolute magnitude $M_B \leq -21$. We find that: 1) the population mix is comparable to that in other ranges of absolute magnitudes; 2) only a small fraction are located in bona fide clusters; 3) the bright galaxy–cluster cross–correlation function is significantly higher on large scales than that measured for fainter galaxies; 4) the correlation length of galaxies brighter than $M_B \sim -20.0$, expressed as a function of the mean interparticle distance, appears to follow the universal dimensionless correlation function found for clusters and radio galaxies; 5) a large fraction of the bright galaxies are in interacting pairs, others show evidence for tidal distortions, while some appear to be surrounded by faint satellite galaxies. We conclude that very luminous optical galaxies differ from the normal population of galaxies both in the clustering and other respects. We speculate that this population is highly biased tracers of mass, being associated to dark halos with masses more comparable to clusters than typical loose groups.

Subject headings: cosmology: observations — dark matter — galaxies:formation — galaxies:fundamental parameters — galaxies: halos — large-scale structure of universe

1. Introduction

Recently, the dependence of the clustering properties of galaxies on luminosity was investigated by Benoist et al. (1996) and Willmer, da Costa & Pellegrini (1998) using volume-limited subsamples drawn from the SSRS2 (da Costa et al. 1994, 1998). Analysis based on the two-point correlation function, in redshift and real space, and counts-in-cells showed that luminous galaxies ($L > L^*$) are more clustered than sub–L* galaxies. The effect is particularly strong for the very luminous galaxies with $M_B \leq -21$ (hereafter VLGs), for which the correlation length approaches that of $R \geq 0$ galaxy clusters.

This result might simply imply that the bright galaxies are preferentially located in clusters. This would be the case, for instance, if these galaxies were primarily cD galaxies, as suggested by Hamilton (1988) in his analysis of the CfA1 catalog. Another possibility is that very bright galaxies are located in loose groups which are expected, on theoretical grounds (e.g. Hamilton & Gott 1988), to have clustering properties intermediate between galaxies and clusters, although no strong observational evidence currently exists in support to this conjecture. Previous attempts to measure the correlation function of loose groups in the CfA and SSRS surveys with a standard algorithm (Huchra & Geller 1982) have led to the conclusion that the amplitude is comparable to –or even lower than– that of galaxies (Jing & Zhang 1988, Maia & da Costa 1990, Ramella et al. 1990). Progress has recently been made by Girardi et al. (1998) analyzing a much larger group catalog derived from CfA2 and SSRS2. These authors show compelling evidence that groups are indeed more clustered than galaxies with a correlation length of the order of 10 h^{-1} Mpc, yielding a relative bias of ≈ 3. They also find evidence that the correlation amplitude increases with luminosity (mass) of the group. Therefore the clustering properties of VLGs might also be explained if VLGs are in very rich groups.

Alternatively, if VLGs are not preferentially in rich groups or poor clusters, an intriguing possibility is that they are associated to massive dark halos and that their large correlation length reflects the relative bias of these halos to the underlying

mass distribution. Semi–analytical models of galaxy evolution such as those of Kauffmann, White & Guiderdoni (1993) are now able to make some predictions about the luminosity function of galaxies in halos of different mass, despite the uncertainties related to the galaxy–dark halo connection. An apparent generic prediction is that each halo would contain a dominant galaxy surrounded by fainter galaxies (Kauffmann, Nusser & Steinmetz 1997). If this is the case, then it would imply that at least some cases of dark matter concentrations can be detected and used to investigate the halo–galaxy connection.

Here we investigate in more detail the characteristics of very luminous galaxies (hereafter VLGs) and compare their clustering properties to those of galaxy clusters. In section 2, we describe the properties of this galaxy population from available data in the literature and investigate the environment in which they reside. In section 3, we examine their distribution relative to the large–scale structures and their auto–correlation and VLG–cluster cross–correlation properties. Our main conclusions are summarized in section 4.

2. Characteristics of Very Luminous Galaxies

2.1. VLG Sample

The VLG sample was drawn from the complete magnitude–limited SSRS2 south catalog, which contains about 3400 galaxies with $m_B(0) \leq 15.5$ (da Costa et al. 1994, 1998), covering the region $b \leq -40^o$ and $-40^o \leq \delta \leq -2.5^o$. Absolute magnitudes were estimated from luminosity distances, assuming $H_0 = 100$ and $q_0 = 0.5$. Velocities were corrected for the motion of Local Group and apparent magnitudes corrected for K–dimming according to the different morphological types. Our VLG sample consists of 113 galaxies with $M_B \leq -21$ ($L > 4L^*$). The redshift distribution for these galaxies is shown in figure 1. We should point out that we excluded from our sample a few cases of strongly interacting systems, or of galaxies nearby bright stars, which had been preliminarily selected in the catalog because their magnitude had

been significantly overestimated (a few VLGs we have included might still be affected by this problem, e.g. VLG 108). Together with the SSRS2 magnitude errors (~ 0.3), there is some uncertainty in our VLG catalog limited at $M \leq -21$, but this has no impact in our conclusions.

2.2. Individual Properties

In order to look for possible explanations for the strong clustering exhibited by the VLGs (see Benoist et al. 1996 and discussion below) we first examine the characteristics of the individual galaxies that comprise this population. For this purpose, we have used the morphological classification as available in the SSRS2 catalog (da Costa et al. 1998) and have searched the NASA Extragalactic Database (NED) for additional information on these galaxies.

Our first finding is that the morphological composition of our VLG sample does not differ from the whole SSRS2 sample. In the bright sample there are 39 (35%) early–type galaxies ($T < 0$) and 74 (65%) late-types. These fractions are consistent, for example, to the 35% of early types found in the $M_B \leq -19.0$ volume–limited subsample, which does not show the enhanced correlation exhibited by the VLGs. It is worth stressing here that morphological classification of these bright galaxies –at least the separation between early and late types– is quite reliable (da Costa et al. 1998).

From the NED we also find that only four galaxies are cDs, which, as mentioned, would be the most natural explanation for the observed strong clustering, as in that case one would be measuring the amplitude of the cluster–cluster correlation.

In order to illustrate the properties of VLG galaxies, we summarize in table 1 the information gathered from the NED for all the VLG galaxies. The table includes: in column (1) the identification number; in column (2) the catalog name; in column (3) the ESO name (Lauberts 1980), whenever available; in columns (4) and (5) the B1950 right ascension and declination; in column (6) the apparent magnitude

as listed in the SSRS2 catalog; in column (7) the B-R color (Lauberts & Valentjin 1989; de Vaucouleurs et al. 1991), whenever available; in column (8) the infrared luminosity at 60μm in solar units (with $H_0 = 100$ km s^{-1} Mpc^{-1}); in column (9) the morphological type either from the SSRS2 catalog or from the literature if more detailed classification is available; in column (10) the heliocentric radial velocity; in column (11) an indication of other properties such as infrared emission (ir), AGN (an), radio source (rs).

We call attention that for the 13 VLG galaxies with radial velocities less than 10,000 km s^{-1}, for which the information should be the most complete, Table 1 shows that late–spirals is the dominant morphological type, often with bars and rings. Moreover, 6 out of the 13 nearby VLGs are also detected by *IRAS*, typically with $L_{IR} \sim 10^{10}$ in solar luminosity units.

For the population as a whole we find that a large number show other interesting characteristics: 1) there are 30 peculiar galaxies (Arp and Madore 1987); 2) 42 galaxies are also in the IRAS Point Source Catalog and/or in the Faint Source Catalog (Moshir et al. 1990); 3) we find an apparent overabundance of barred and ring galaxies. Out of 26 galaxies with detailed morphological information 13 contain bar/ring structure, while 7 are intermediate cases; 4) 14 galaxies are radio–sources; 5) only 5 galaxies are AGNs (3 Seyferts and 2 Liners; Maia et al. 1996, NED), showing that the VLGs have luminosities that exceed some nearby AGN–like galaxies.

In the VLG sample, we find that 54 are included in the ESO catalog for which B and R magnitudes are available. In figure 2 we compare the color distribution of this subsample with that of the $M \leq -19$ sample, renormalized to the total number of VLGs. While VLGs have a slightly larger spread in color, the 2 distributions are similar.

The above results indicate that the sample of SSRS2 galaxies drawn on the basis of their large blue luminosity is "special" relative to galaxies of lower luminosity both in terms of their internal characteristics as well as their clustering properties.

2.3. Relation VLGs–Clusters

As suggested earlier, an obvious explanation for the large clustering strength of bright galaxies would be that we are simply picking up cluster members, thus measuring the amplitude of the cluster–cluster correlation function (e.g., Hamilton 1988). Although, as previously mentioned, the VLGs are not cDs, they could still reside preferentially in rich clusters. Even though the volume covered by the SSRS2 does not include very rich clusters, we have examined the ACO catalog (Abell, Corwin & Olowin 1989), which can be considered complete out to a distance of 200 h^{-1} Mpc, searching for VLGs with a projected separation less than 1.5 h^{-1} Mpc from cataloged $R \geq 0$ cluster centers and a radial velocity difference less than 1200 km s^{-1} with respect to the cluster mean velocity. We find only 12 galaxies which can be considered as candidate cluster members. We have also extended our search to the poor clusters in the volume, using the ACO supplementary list and the Edinburgh–Durham cluster catalog (Nichol et al. 1992), with a measured redshift. In this search we include clusters beyond the redshift limit of the bright galaxy sample. Using the same criteria as that used for the rich clusters we find 7 additional galaxies which could be associated with known poor clusters. Therefore, at best only 19 VLGs may reside in previously known rich and poor clusters, out of which 16 are early–type galaxies. This result confirms that very luminous galaxies are not preferentially in $R \geq 0$ or rich clusters.

In figure 3 we show the spatial distribution of the VLGs and the ACO clusters in the region, in the declination range $-40^o \leq \delta \leq -2.5^o$. In order to have a preliminary understanding of the VLG location relative to the large–scale structures, we have resorted to a simple percolation analysis. Adopting a percolation parameter $s = 0.5$, which corresponds to a search radius $r_s = 12.6$ h^{-1} Mpc and to a space density enhancement of 1.9, we have detected 7 structures with more than 3 members, containing about 42% of the bright galaxies. Among them there are two main structures. One is at a mean redshift $z \sim 0.055$. It consists of 20 members, out of which only 5 are possible members of rich clusters, all belonging to the Pisces–Cetus

region. The second association, with 9 members, is at a mean redshift $z \sim 0.054$, and none of its galaxies are found in rich clusters. Comparing their coordinates with maps of the large–scale structures, based on the ACO cluster distribution (Tully 1986, Tully et al. 1992), both these associations correspond to dense regions of the Pisces–Cetus Supercluster.

As a final note, we point out that eliminating the VLGs which may be in clusters does not significantly change the correlation function of the sample (see section 3).

2.4. Environment of VLGs

As we have seen, there is no compelling evidence that VLGs reside preferentially in clusters of galaxies. However, this does not fully answer the more general question about the type of environment in which these galaxies reside. Unfortunately, this is not an easy task because the majority of the VLGs are at large distances, where fainter neighbors would not be included in the SSRS2 sample because of its relative bright magnitude limit. On the other hand, the use of radial velocity databases is dangerous because of their unknown incompleteness. Therefore, we can only attempt a preliminary investigation of the question.

First of all, we have used the catalog of groups recently identified within the SSRS2 sample by Girardi et al. (1998). This catalog contains groups with mean positions out to 12,000 km s^{-1}, but includes members out to 15,000 km s^{-1}. With this procedure we should in principle be able to detect groups including nearby VLGs: at 12,000 km s^{-1} we can detect a group with a VLG and 2 companions brighter than ~ -19.9 or even fainter at smaller distance (there are 27 VLGs within $V \leq 12,000$km s^{-1}, and 45 VLGs within $V \leq 15,000$km s^{-1}). We find 14 VLGs in 13 loose groups; of these, 3 are in known clusters (and have already been included in the 19 mentioned above), 3 are in known groups –2 in Hickson compact groups 12 and 91 (see Hickson 1982), one in an Arp–Madore group–, and 8 are in new groups, hereafter referred as SSRS2 groups. The richness of the 8 loose groups (i.e. those which are

not associated with rich clusters or compact groups) is quite low; 4 groups have 3 members, 2 groups have 5 members, and 1 has 7 members, including 2 VLGs.

The identification of groups in the total VLG sample is more difficult, and is necessarily incomplete. Using the whole SSRS2 catalog, we have selected galaxies with a projected separation of 1.5 h^{-1} Mpc and within 1200 km/s from a very luminous galaxy. This is not a real group finding algorithm but it suffices for our purposes. In this way we have found 22 galaxies in group candidates with at least 3 members (including the VLG); most of them are known systems we have already taken into account: only 4 are new candidate systems, which have not been identified in the catalogs of clusters or groups.

In addition, we have also searched VLG galaxies in the compact group catalog by Barton et al. (1996). We find two systems which include nearby VLGs: their no.81 (which corresponds to HCG91) and no.88, which corresponds to the core of A4038. Both these cases are also in the SSRS2 group catalog (Girardi et al. 1998).

However, as we have stated the number of systems we can identify is biased by the relatively bright magnitude limit of the SSRS2, which could lead us to miss fainter galaxies. Therefore, we have also searched the literature (e.g., NED) to identify known systems of galaxies from binaries to groups. Of course, one should be cautious in interpreting the results because of the incompleteness of these data sets. From this search we find for example other 2 VLGs in southern compact groups from the automated catalog by Prandoni, Iovino & MacGillivray (1994), 13 pairs and 4 triplets (2 pairs belonging to systems previously found).

Combining all the above results we may conclude that out of the 113 VLGs we have identified 12 in rich clusters, 7 in poor clusters, 24 in generic groups with at least 3 members and 11 in interacting pairs or possible binaries, for a total of 54 VLGs in some type of galaxy association. In table 2, we list all VLGs that have been identified with some system ranging from binaries to rich cluster. In column (1) we give the identification number, in column (2) the catalog name, in column (3) the system type (binary, triplet, group, poor cluster, rich cluster), with an eventual

note on the VLG (cD, Seyfert, interacting), in column (4) the system name (ACO or EDCC name for clusters, Hickson no. for compact groups, Arp–Madore, ESO or Vorontsov–Velyaminov no. for the others, indicating simply SSRS2 for groups listed only in the Girardi et al. (1998) catalog or for candidate groups found by percolation in this work), in column (5) the velocity dispersion of the system as measured from the SSRS2, in column (6) the number of galaxies used to estimate the velocity dispersion. As apparent from table 2, there is generally only one VLG per system. Two exceptions are A4008, which contains 4 VLGs, and the SSRS2 loose group which includes VLG 37 and VLG 38.

From the above identified systems, extended X–ray emission, consistent with the VLG position, was detected by ROSAT for galaxies in A4038 and S0141. The object VLG 86, which is a Seyfert 1, was detected as a point source in X by ROSAT (Böhringer, private communication). At the position of VLG 65, ROSAT detected also X–ray extended emission typical of a group or cluster. This is an interesting object, because even if it is relatively nearby ($z \sim 0.03$), it is not found in any cluster or group catalog. We identified it in the SSRS2 through percolation as a group of 3 galaxies (including the VLG) and in a note in the MCG (Vorontsov–Velyaminov & Arhipova 1968) referring to this galaxy we read: "Here is the main member of a cluster, located at its end. The other members are of type E and S0 and are much fainter, but visibly spiral and [16-17 mag]."

The remaining 59 VLGs have not been associated to any particular known system. However, after eye–examination of these galaxies using the Digitized Sky Survey (DSS), we estimate that only about 16 galaxies might be isolated. Among the other galaxies, about 20 have at least a companion, and show clear distorted morphologies and/or evidence of interaction, 5 appear to be surrounded by faint satellites and 18 might be in groups (often with members much fainter than the VLG, as in the case of VLG 65), although usually no redshift information is available. This is illustrated in figure 4 for four such cases: VLG 24 is surely not a typical case, being an elliptical galaxy presumably at the center of a group; VLG 41 is probably

a member of a triplet/small group of galaxies; VLG 44 is clearly interacting with a large, irregular galaxy; VLG 77 is a Liner, probably interacting with a small satellite.

Another evidence that VLGs are unlikely to be isolated systems, comes from the examination of the small subsample of nearby VLGs ($V < 10,000$ km s^{-1}, a distance at which a typical $M^* \sim -19.5$ galaxy is brighter than m=15.5 and is therefore included in the SSRS2). Images for these galaxies are shown in figure 5. Most of them are not isolated. For example, VLG 20 does not belong to any known cluster or group, but we have identified around it (through the percolation technique previously described) other 7 fainter SSRS2 galaxies, giving a velocity dispersion of ~ 900 km/s (see table 2). Close inspection of the images for the only three nearby VLGs, VLG 27, VLG 29, and VLG 61, which, from our analysis, would be classified as "isolated", show instead that they have companions: VLG 27, a peculiar barred late–type spiral, seems to be surrounded by satellites very close it as well as brighter galaxies at larger distances; VLG 29 (listed in the Arp–Madore catalog of Southern Peculiar galaxies) has a disk galaxy as a probable companion; VLG 61 has many nearby satellites.

From the above evidence we can conclude that VLGs are generally found in environments of high local galaxy density, but not in rich clusters. Moreover, both eye-inspection of the images of nearby VLGs and their surroundings (within 1 h^{-1} Mpc at the VLG redshift), and the number of members in SSRS2 loose groups found to include a VLG, show no evidence that VLGs are preferentially in rich groups.

3. Large–scale Distribution

3.1. Correlation function

The original reason for focusing our attention in the sample of VLGs was the strong bias showed by this population relative to fainter galaxies (Benoist et al. 1996). This is illustrated in figure 6 where we show the correlation function for galaxies brighter than $M = -21$ within 168 h^{-1} Mpc. For comparison, we also show the fit to the cluster correlation function (Cappi & Maurogordato 1992) determined for $R \geq 0$

ACO clusters (Abell et al. 1989), $\xi_{cc} = (s/19.5)^{-1.8}$.

We remind that the correlation function of clusters depends on their richness (see Mann et al. 1993, Croft et al. 1997) For $R \geq 1$ clusters Peacock & West (1992) find 21 ± 1.3 h^{-1} Mpc, but some authors claim that the correlation amplitude of Abell clusters is amplified by projection effects, and give an estimate of ~ 14 h^{-1} Mpc for APM clusters (Dalton et al. 1994), which would then be comparable to the the value found for VLGs.

From the figure we see that very bright galaxies have a correlation length comparable to that measured for poor clusters. The correlation length of $M_B \leq -21$ galaxies is $r_0 = 16 \pm 2$ h^{-1} Mpc, while the zero–crossing of the correlation function is beyond ~ 40 h^{-1} Mpc.

In Benoist et al. (1996), it was possible to prove that the dependence of the correlation amplitude on luminosity was real, by considering different luminosity-limited samples within the same volume. Unfortunately, the same direct test is not possible for the $M \leq -21$ sample, because of the relatively small number of galaxies. Therefore, one might argue that the large correlation amplitude is due to sampling effects, caused either by virialized systems in the volume considered or by fluctuations of the mean background density, on scales comparable to the sample size; in fact, examining figure 3, where we show the spatial distribution of the bright galaxies, in the declination range $-40^o \leq \delta \leq -2.5^o$, at least two concentrations can be seen at large distances, at $\alpha \sim 23^h 30^m$ and at $\alpha \sim 1^h$, corresponding to large–scale structures seen in redshift space, as discussed in section 2.3.

There are, however, two lines of argument showing that our results are not a consequence of sampling effects. The first, based on circumstantial evidences, is that: 1) as we have seen the bright galaxies seem to have peculiar characteristics; 2) their clustering properties extend a trend already visible for fainter galaxies (see figure 5 of Benoist et al. 1996); 3) similar results were found by Hamilton (1988) in his analysis of the CfA1 catalog, i.e. probing a different region of the sky, and by Park et al. (1994) from the analysis of the CfA2 power spectrum. A more direct argument is

based on the following facts: 1) the correlation amplitude does not vary significantly if one considers sub-samples defined by removing the main concentrations; 2) by using a smaller volume; 3) or by splitting the sample into two ranges of right ascension, probing different structures.

Therefore, the large correlation amplitude of very bright galaxies is probably a genuine property. We also point out that the amplitude of the correlation is not affected by removing galaxies identified in the previous section as possibly belonging to rich clusters. It is also important to emphasize that although a large fraction of the VLGs may be in systems of galaxies, currently there is no evidence for the amplitude of the SSRS2 group–group correlation function (Girardi et al. 1998) to be as large as that obtained for the VLGs. On the basis of the evidence presented in the previous section, VLGs are not preferentially in clusters, and probably not in rich groups (this last claim is obviously based on our analysis of nearby VLGs). We therefore suggest that VLGs are a highly biased population, possibly associated with dark halos with masses comparable to clusters.

3.2. VLG–Cluster cross–correlation function

An additional evidence in support of the above conjecture is the VLG-cluster cross-correlation function shown in figure 7a. Despite the small number of clusters (we have found only 28 $R \geq 0$ ACO clusters within the volume considered), we find $\xi_{gc} = (r/16 \pm 2)^{-1.7\pm0.4}$. For comparison we also show the galaxy–cluster cross-correlation determined by Lilje & Efstathiou (1988), $\xi_{gc} = (r/8.8)^{-2.2}$, based on the Lick counts of galaxies. While the slopes differ at the 1σ level, r_0 for bright galaxies is significantly larger, by a factor of about 2, even though the uncertainties are large (for example, Seldner & Peebles 1977 find a significantly larger amplitude for the galaxy–cluster cross–correlation than Lilje & Efstathiou 1988).

An estimate of the relative bias between the VLGs and clusters can be computed from the ratio $\alpha = J_3(gc)/J_3(cc)$, where $J_3(s) = \int_0^s x^2 \xi(s) ds$, $J_3(gc)$ is the integral of

the galaxy–cluster cross–correlation, and $J_3(cc)$ is the integral of the cluster correlation function (see Alimi et al. 1988). The dependence of α on separation s is shown in figure 7b from which we determine that galaxies with $M_B \leq -21$ are less clustered by a factor of $b_{21}/b_{cc} \sim 0.8$ relative to clusters, which appears to be constant in the range 5–50 h^{-1} Mpc. This high bias is comparable to that measured for radio–galaxies (Peacock & Nicholson 1991, Mo, Peacock & Xia 1993, Loan, Wall & Lahav 1997), and which has been explained as being due to the fact that radio–galaxies reside preferentially in clusters. However, as we have seen the VLGs are neither radio-sources nor are located in rich clusters. Furthermore, while radio–galaxies are representative of early–type galaxies, our results show that late–type galaxies also contribute to the effect.

3.3. Dimensionless Correlation Function

In order to further compare the clustering properties of the bright galaxies with those of clusters and other systems we use the scaling relation proposed by Szalay & Schramm (1985), who noted that the cluster correlation amplitude increases with their mean spatial separation. As discussed by Bahcall & West (1992) the "universal dimensionless correlation function"

$$A_i = r_0^{\gamma} \sim (\alpha d_i)^{1.8} \tag{1}$$

where $d_i = n^{-1/3}$ is the mean inter-particle distance and $\alpha \sim 0.5$, seems to hold for a number of systems up to superclusters, but not for normal galaxies. However, our results suggest that in this respect $M \leq -21$ galaxies should differ from typical M^* galaxies, as their mean density is much lower ($n \sim 6 \times 10^{-5}$ h^3 Mpc^{-3} in the SSRS2), and they are more strongly correlated.

In figure 8 we show the correlation length r_0 for different volume-limited subsamples of the SSRS2, as a function of their mean inter-particle distance $d_i = n^{-1/3}$, where n is the mean density of the subsample. Filled hexagons refer

to SSRS2 volume–limited subsamples: each subsample includes galaxies in a one magnitude range, from $-17 \leq M_B \leq -16$ (lowest point) to $-21 \leq M_B \leq -22$, with a step of 0.5 magnitudes (for details, see Benoist et al. 1996).

Recalling that for the SSRS2, $M_B{}^* \sim -19.5$, it is clear from figure 8 that faint galaxies ($L < L^*$) show a significant departure from the relation, as one could expect *a priori* (see Peebles 1993), whereas brighter galaxies ($L > L^*$) approach it asymptotically, which is a much less trivial result. This is in fact the first time that it is possible to cover a sufficiently large range of luminosities to allow a comparison of the scaling relation of galaxies and clusters. Galaxies brighter than $M_B = -20$ approximately follow a scaling relation with $\alpha \sim 0.6$, slightly higher but, given the uncertainties, consistent with the value of $\alpha \sim 0.4$ determined by Bahcall & West (1992) for a variety of systems.

Szalay & Schramm (1985; see also Luo & Schramm 1992) argued that the universal dimensionless correlation function followed by clusters could be explained if structures formed by a truly scale–invariant process, leading to a fractal–like distribution, while the observed deviation of galaxies from this relation could be explained by non–linear gravitational clustering at small scales. We take a different point of view and argue that the observed behavior is a clear evidence for biasing. First, because the results of Benoist et al. (1996) show that the correlation length does not depend on the depth of the sample as expected for fractals (see also Cappi et al. 1998). Second, the clear trend of very bright galaxies to lie close to the relation obeyed by clusters suggests a continuity of the clustering properties of bright galaxies and clusters. This would be a natural consequence of different biasing amplitudes for systems with different mass scales (e.g. Peebles 1993).

Our results show that the universal dimensionless correlation function can be explained as a consequence of the increase of the correlation amplitude with the mass (and therefore rarity) of the system, as expected if the distribution of luminous objects is biased with respect to the underlying matter distribution (Kaiser 1984; Bardeen et al. 1986). The $r_0 - d$ relation should thus be regarded as an empirical tool to compare

the biasing of different classes of objects (see for example the more rigorous approach of Bernardeau & Schaeffer 1992).

4. Discussion and Conclusions

From our analysis of the SSRS2, we have shown that very luminous galaxies ($M_B \leq -21$ or $L \gtrsim 4L^*$) are strongly correlated ($r_0 = 16 \pm 2$ h^{-1} Mpc) and the zero–crossing of the correlation function is beyond 40 h^{-1} Mpc. Furthermore, we find that luminous galaxies with $L > L^*$ appear to follow a universal dimensionless correlation similar to that of galaxy clusters, which implies a common biasing mechanism.

Independently of their clustering properties we have found that $L \gtrsim 4L^*$ galaxies seem to form a special population. They are observed in locally high galaxy density regions, but are not predominantly in clusters nor apparently in rich groups. Instead, they tend to be in interacting pairs, some exhibit tidal distortions, while others have faint satellite galaxies in their surroundings. VLGs resemble another class of galaxies: the ultra–luminous infrared galaxies, where the high luminosity appears to be triggered by interaction with other galaxies (e.g. Melnick & Mirabel 1990; Clements et al. 1996; Duc et al. 1997). Is there an evolutionary link between these systems? Is it possible that a massive galaxy undergoes a period of intense star formation, shrouded by dust, and becomes finally visible as a VLG in the optical? A simple test of this hypothesis will be possible with a large and complete sample of infrared ultra–luminous galaxies (which should have a higher correlation amplitude than normal IRAS galaxies).

The fact that the brightest galaxies in the sample are not found in clusters but have clustering properties similar to $R = 0$ clusters suggests that they may reside within massive dark halos. We cannot completely exclude that VLGs are preferentially in rich groups, which should be more strongly clustered than loose groups as a whole, because most VLGs are at large distances, where fainter members are not included in

the SSRS2. However, as discussed above, at the present time we find no evidence for that, as most groups containing a nearby VLG are of low richness. While many VLGs have companions, the total luminosity of these systems appear to be far less than that typical of clusters. If this is the case, it implies that there are large variations in the M/L ratio and/or in the luminosity function of galaxies forming in dark halos. Further observational work is clearly needed to fully characterize this galaxy population. Clarifying the nature of these objects may contribute to our understanding of the halo–galaxy connection and the mechanisms responsible for galaxy evolution.

This research has made use of the NASA / IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. The Digitized Sky Surveys were produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope. The plates were processed into the present compressed digital form with the permission of these institutions. LNdC thanks his SSRS2 collaborators for allowing to use the data prior to publication and the hospitality of the Institut d'Astrophysique and the Observatoire de Paris–Meudon. AC thanks the hospitality of the Observatoire de Nice. We wish also to thank H. Böhringer for his list of X–ray detection in the fields of very luminous galaxies, and the referee M. Ramella for his useful comments.

REFERENCES

Abell G.O., Corwin H.G., Olowin R.P., 1989, ApJS 70, 1 (ACO)

Alimi J., Valls–Gabaud D., Blanchard A., 1988, A&A 206, L11

Arp H.C., Madore B.F., 1987, *Catalogue of Southern Peculiar Galaxies and Associations (2 volumes)*, Cambridge University Press

Bahcall N.A., Chokshi A., 1992, ApJ 385, L33

Bahcall N.A., West M.J., 1992, ApJ 392, 419

Bardeen J.M., Bond J.R., Kaiser M., Szalay A.S., 1986, ApJ 304, 15

Barton E., Geller M.J., Ramella M., Marzke R.O., da Costa L.N., 1996, AJ 112, 871

Benoist C., Maurogordato S., da Costa L.N., Cappi A., Schaeffer R., 1996, ApJ 472, 452

Bernardeau F., Schaeffer R. 1992, A&A 255, 1

Cappi A., da Costa L.N., Benoist C., Maurogordato S. 1998, A&A, submitted

Cappi A., Maurogordato S., 1992, A&A 259, 423

Clements D.L., Sutherland W.J., McMahon R.G., Saunders W., 1996, MNRAS 279, 477

Croft R.A.C., Dalton G.B., Efstathiou G., Sutherland W.J., Maddox S.J., 1997, MNRAS 291, 305

da Costa L.N., Geller M.J., Pellegrini P.S., Latham D.W., Fairall A.P., Marzke R.O., Willmer C.N.A., Huchra J.P., Calderon J.H., Ramella M., Kurtz J., 1994, ApJ 424, L1

da Costa L.N. et al., 1998, AJ, in press

Dalton G.B., Croft R.A.C., Efstathiou G., Sutherland W.J., Maddox S.J., Davis M., 1994, MNRAS 271, L47

de Vaucouleurs G., de Vaucouleurs A., Corwin Jr. H.G., Buta R. J., Paturel G., Fouqué P., 1991, *Third Reference Catalogue of Bright Galaxies*

Duc, P.A., Mirabel I.F., Maza J., 1997, A&AS 124, 533

Girardi M. et al. 1998, in preparation

Hamilton A.J.S., 1988, ApJ 331, L59

Hamilton A.J.S., Gott J.R. III, 1988, ApJ 331, 641

Hickson P., 1982, ApJ 255, 382

Huchra J.P., Geller M.J., 1982, ApJ 257, 423

Jing Y., Zhang J., 1988, A&A 190, L21

Kaiser N. 1984, ApJ 284, L9

Kauffmann G., Nusser A., Steinmetz M., 1997, MNRAS 286, 795

Kauffmann G., White S.D.M., Guiderdoni B., 1993, MNRAS 264, 201

Lauberts A., 1982, *The ESO/UPPSALA survey of the ESO(B) atlas*, Garching bei München, European Southern Observatory

Lauberts A., Valentijn E.A., 1989, *The surface photometry catalogue of the ESO–UPPSALA galaxies*, Garching bei München, European Southern Observatory

Lilje P.B., Efstathiou G., 1988, MNRAS 231, 635

Loan A.J., Wall J.V., Lahav O., 1997, MNRAS 286, 994

Luo X., Schramm D.N., 1992, Science 256, 513

Maia M.A.G., da Costa L.N., 1990, ApJ 349, 477

Maia M.A.G., da Costa L.N., Willmer C.N.A., Rite C., 1996, A&AS 117,487

Mann R.G., Heavens A.F., Peacock J.A., 1993, MNRAS 263, 798

Melnick J., Mirabel I.F., 1990, A&A 231, L19

Mo H.J., Peacock J.A., Xia X.Y., 1993, MNRAS 260, 121

Moshir M., Kopan G., Conrow T., McCallon H., Hacking P., Gregorich D., Rohrbach G., Melynk M., Rice W., Fullmer L., et al., 1990, *Infrared Astronomical Satellite Catalogs, The Faint Source Catalog, version 2.0*

Nichol R.C., Collins C.A., Guzzo L., Lumsden S.L., 1992, MNRAS 255, 21

Park C., Vogeley M.S., Geller M.J., Huchra J.P., 1994, ApJ 431, 569

Peacock J.A., Nicholson D., 1991, MNRAS 253, 307

Peacock J.A., West M.J., 1992, MNRAS 259, 494

Peebles P.J.E., 1993, in *Principles of Physical Cosmology*, Princeton University Press

Postman M., Huchra J.P., Geller M.J., 1992, ApJ 384, 407

Prandoni I., Iovino A., MacGillivray H.T., 1994, AJ 107, 1235

Ramella M., Geller M.J., Huchra J.P., 1990, ApJ 353, 51

Seldner M., Peebles P.J.E., 1977, ApJ 215, 703

Szalay A., Schramm D., 1985, Nature 314, 718

Tully R.B., 1986, ApJ 303, 25

Tully R.B., Scaramella R., Vettolani G., Zamorani G., 1992, ApJ 388, 9

Vorontsov–Velyaminov B., 1977, A&AS 28, 1

Vorontsov–Velyaminov B., Arhipova V.P., *Morphological Catalogue of Galaxies*, 1968, Volume IV, Moscow: Moscow State university

Willmer C.N.A., da Costa L.N., Pellegrini P.S., 1998, AJ, in press

---

This manuscript was prepared with the AAS LaTeX macros v4.0.

Table captions

Table 1 — List of the VLGs galaxies ($M_B \leq -21$) in the SSRS2; ir=detected by IRAS, rs=radiosource, an=AGN.

Table 2 — List of VLGs in known systems of galaxies.

Figure captions

Fig. 1.— Redshift distribution of VLG galaxies.

Fig. 2.— Histogram Comparison of the $B-R$ color distribution of VLGs (solid line) and $M \leq -19$ galaxies (dashed line; renormalized to the total no. of VLGs).

Fig. 3.— Spatial distribution of VLGs and ACO cluster. The cone diagram includes the $M_B \leq -21$ galaxy subsample (stars), and the ACO clusters in the same b_{II} and δ limits up to a redshift $z = 0.07$ (triangles).

Fig. 4.— Four examples of distant VLGs with distorted morphology, faint satellite galaxies and/or companions or IG. Fields are about 11×11 arcminutes (corresponding to $\sim 2 \times 2$ h^{-1} Mpc at the typical distances of these galaxies).

Fig. 5.— Mosaic showing the images of the 13 nearby VLGs ($M \leq -21$ and $V \leq 10000$ km/s). Fields are 15×15 square arcminutes.

Fig. 6.— Comparison of the VLG and cluster autocorrelation functions. Filled hexagons: volume–limited sample, $D_{lim} = 168.5$ h^{-1} Mpc, for galaxies with $M_B \leq -21$. Solid straight line: fit for $R \geq 0$ ACO clusters (Cappi & Maurogordato 1992).

Fig. 7.— a) Galaxy–cluster cross–correlation function for galaxies brighter than $M_B = -21$ (subsample limited at $D_{lim} = 168.5$ h^{-1} Mpc) The solid line gives the best fit to the data. The dashed line is the fit to the standard galaxy–cluster cross–correlation by Lilje & Efstathiou (1988). b) The ratio $\alpha = J_3(gc)/J_3(cc)$ as a function of separation.

Fig. 8.— The correlation length r_0 as a function of the mean interparticle distance $d_i = n^{-1/3}$. Filled hexagons: galaxy subsamples of the SSRS2 from $M_B = -16$ to $M_B = -21$. Asterisk: CfA2+SSRS2 groups from Girardi et al. (1998). Filled triangle: radiogalaxies (from Bahcall & Chokshi 1992). Filled squares: 2 northern and southern ACO cluster subsamples with $R \geq 0$ as defined in Cappi & Maurogordato 1992. Open hexagon: $R \geq 1$ Abell clusters. Star: very rich Abell clusters ($R \geq 2$; see Bahcall &

West 1992). Lines represent the relation $A_i \sim (\alpha d_i)^{1.8}$. Solid line: $\alpha = 0.4$; dotted line: $\alpha = 0.5$; dashed line: $\alpha = 0.6$.

Fig. 1.—



Fig. 2.—



Fig. 3.—


REDSHIFT
RA
21 0 0
5 0 0
0.01
0.02
0.03
0.04
0.05
0.06
0.07
DEC FROM-40 0 0 TO -2 30 0

Fig. 6.—



Fig. 7.—



Fig. 8.—



TABLE 1.

No.	Catalogue Name	Other Name	RA	Dec	m_B	B-R	$\log_{10}L_{60}$	Type (km/s)	$V_\odot$	Notes
1	VLG 0006-3408	ESO 349- G 032	00 06 02.2	-34 08 11	13.19	1.51	9.62	SAB(rs)b	6813	ir
2	VLG 0010-2429	ESO 472- G 020	00 10 30.7	-24 29 34	14.00	2.24	10.00	SA(r)c	10269	ir
3	VLG 0012-2458	ESO 473- G 003	00 12 33.8	-24 58 11	15.13	0.73	10.27	Sb	16867	ir
4	VLG 0036-1541		00 36 08.0	-15 41 39	14.87		10.41	5	16352	ir
5	VLG 0039-1708		00 39 15.2	-17 08 05	14.47			SB(s)d	16239	
6	VLG 0039-0934		00 39 18.6	-09 34 38	13.98			cD/SB(r)	16536	rs
7	VLG 0041-2522	ESO 474- G 020	00 41 46.2	-25 22 58	15.13	1.12		Sb	15921	
8	VLG 0042-2839		00 42 22.9	-28 39 47	14.98		10.32	SA(r)a	15061	ir
9	VLG 0044-2438	ESO 474- G 026	00 44 39.8	-24 38 37	15.19	0.97	10.52	SA(s)a	16246	ir
10	VLG 0044-0705		00 44 51.3	-07 05 03	15.21			-5	16843	
11	VLG 0044-1133		00 44 53.5	-11 33 43	15.00			6	17157	
12	VLG 0046-2350	ESO 474- G 031	00 46 15.5	-23 50 03	15.36	1.60		S0	16987	
13	VLG 0047-1457		00 47 30.9	-14 57 16	15.12			-5	16086	
14	VLG 0049-2735		00 49 48.9	-27 35 59	14.50			Sbc	12045	
15	VLG 0050-2314	ESO 474- G 036	00 50 18.7	-23 14 47	15.20	1.44		S0:	15846	
16	VLG 0052-3535	ESO 351- G 021	00 52 37.1	-35 35 33	15.21	1.73		SA0-	17254	
17	VLG 0053-1015		00 53 20.9	-10 15 21	15.14			-2	16352	
18	VLG 0053-3740	ESO 295-IG 022	00 53 24.2	-37 40 43	15.23			E	16657	
19	VLG 0053-1337		00 53 34.3	-13 37 43	15.18			5	16710	
20	VLG 0054-1011		00 54 11.6	-10 11 05	12.50		9.94	SAB(r)c	5666	
21	VLG 0056-1644		00 56 22.4	-16 44 19	14.70			-2	16226	
22	VLG 0056-2834		00 56 27.6	-28 34 23	15.31			S0	17079	
23	VLG 0057-1415		00 57 29.5	-14 15 50	15.20			-2	15972	
24	VLG 0057-2145	ESO 541- G 007	00 57 34.9	-21 45 28	15.09	1.59		E-S0	16805	
25	VLG 0057-1534		00 57 46.8	-15 34 06	15.28			-2	16235	rs
26	VLG 0100-2209	ESO 541- G 013	01 00 15.1	-22 09 02	14.84	1.60		cD;E+3pec	17073	rs
27	VLG 0101-3523	ESO 352- G 001	01 01 57.8	-35 23 21	14.04	1.08	10.36	SB(r)bc pec	9991	ir
28	VLG 0103-1916		01 03 37.1	-19 16 19	15.19			-2	16194	
29	VLG 0103-3026	ESO 412- G 00	01 03 49.3	-30 26 43	13.72	1.10	10.44	SB(r)c	9620	ir
30	VLG 0106-1606		01 06 19.4	-16 06 38	14.46		9.74	SA(rs)a	12073	ir

arXiv:astro-ph/9803033v1 4 Mar 1998

TABLE 1. (continued)

No.	Catalogue Name	Other Name	RA	Dec	m_B	B-R	$\log_{10}L_{60}$	Type (km/s)	$V_\odot$	Notes
31	VLG 0106-1540		01 06 22.6	-15 40 24	14.36			D	15967	
32	VLG 0111-3200	ESO 412- G 018	01 11 26.4	-32 00 43	12.59	1.57		SAB(rs)0-	5803	
33	VLG 0116-2547		01 16 49.8	-25 47 35	14.62		10.75	SA(s)a	16188	ir
34	VLG 0117-0530		01 17 58.9	-05 30 59	14.96			5	17040	
35	VLG 0125-0456		01 25 01.7	-04 56 31	14.96			S0	14407	
36	VLG 0130-1504		01 30 24.8	-15 04 16	14.33		10.05	5	11387	ir
37	VLG 0131-3644	ESO 353- G 015	01 31 43.5	-36 44 57	13.16	1.57	10.25	SA0+pec sp	8985	
38	VLG 0132-3644	ESO 353- G 023	01 32 52.4	-36 44 44	13.88	1.64		E+:	8955	
39	VLG 0134-0927		01 34 46.2	-09 27 08	14.33			SA0-	12394	rs
40	VLG 0136-2849		01 36 41.1	-28 49 33	14.64			S	12628	
41	VLG 0137-0929		01 37 27.0	-09 29 43	14.50			SB(r)0?	12741	
42	VLG 0137-1506		01 37 33.4	-15 06 56	14.93		10.36	5	14973	ir
43	VLG 0137-2812	ESO 413- G 015	01 37 38.2	-28 12 34	15.28	0.99		Sb	16825	
44	VLG 0146-1238		01 46 05.0	-12 38 09	14.92		10.27	5	14301	ir
45	VLG 0147-2907		01 47 57.4	-29 07 10	14.95		10.01	S	17257	ir
46	VLG 0148-0418		01 48 47.5	-04 18 06	14.23		10.33	5	10607	
47	VLG 0150-1901	ESO 543- G 022	01 50 16.8	-19 01 39	14.92			S?	14580	
48	VLG 0152-2823	ESO 414-IG 009	01 52 32.0	-28 23 38	15.17	1.12	10.37	4	16945	ir
49	VLG 0154-2017	ESO 543- G 026	01 54 10.8	-20 17 33	15.08		9.84	Sb:	15252	ir
50	VLG 0158-3327	ESO 354- G 029	01 58 58.6	-33 27 02	14.09	1.20	9.83	Sc	10644	ir
51	VLG 0159-2200	ESO 544- G 007	01 59 56.6	-22 00 12	14.47	1.45	9.89	S?	13161	ir
52	VLG 0210-2242	ESO 478- G 013	02 10 11.5	-22 42 20	14.13	0.68	10.44	SB(rs)c?	12356	ir
53	VLG 0210-2641		02 10 33.3	-26 41 35	15.16			S0	17067	
54	VLG 0212-2504	ESO 478- G 015	02 12 32.5	-25 04 59	14.41	1.17		R('2SB(s)b	11394	
55	VLG 0213-2416		02 13 57.5	-24 16 37	15.37		10.05	S	17483	ir
56	VLG 0215-2336	ESO 478- G 022	02 15 36.8	-23 36 53	14.38	1.13	10.19	Sa?	11332	ir
57	VLG 0222-1922	ESO 545- G 013	02 22 20.5	-19 22 02	13.59	1.15	10.33	SAB(s)c Sy1.	10132	ir,an
58	VLG 0223-2411	ESO 479- G 003	02 23 44.6	-24 11 07	15.24	1.80		S0	16266	
59	VLG 0224-2338		02 24 19.8	-23 38 28	14.76			S0	15982	
60	VLG 0227-1715		02 27 17.4	-17 15 57	15.09			3	15872	rs

TABLE 1. (continued)

No.	Catalogue Name	Other Name	RA	Dec	m_B	B-R	$\log_{10}L_{60}$	Type (km/s)	$V_\odot$	Notes
61	VLG 0228-0309		02 28 11.1	-03 09 38	12.73		10.39	SB(rs)c	5738	
62	VLG 0229-2006	ESO 545- G 030	02 29 27.8	-20 06 18	13.82	1.61		SA0 pec	10165	
63	VLG 0231-2115	ESO 545- G 036	02 31 38.0	-21 15 43	15.36	1.55		R'?SA(s)a	17137	
64	VLG 0233-2544	ESO 479- G 015	02 33 07.8	-25 44 43	15.10	1.34		SB(rs)bc	16799	
65	VLG 0253-1424		02 53 22.1	-14 24 34	13.76			SA0-:	8703	rs
66	VLG 0304-0955		03 04 03.1	-09 55 24	14.20			SA0- pec:	10239	
67	VLG 0318-3842	ESO 301-IG 006	03 18 39.0	-38 42 52	15.08	1.40		1	16068	
68	VLG 0322-0621		03 22 43.5	-06 21 22	13.97		10.57	S Sy1	10107	ir,an
69	VLG 0327-2856	ESO 418- G 007	03 27 52.0	-28 56 38	13.95	1.49	9.86	SB(s)b pec	11052	ir
70	VLG 0337-3341	ESO 358-IG 047	03 37 14.3	-33 41 32	14.84	0.56	10.61	S pec	16008	ir
71	VLG 0340-3811	ESO 301-IG 025	03 40 09.3	-38 11 28	14.76	0.91	10.45	SB(s)I-II	13542	ir
72	VLG 0355-1921	ESO 549- G 042	03 55 24.5	-19 21 34	14.36	1.64		SB(r)b	11449	
73	VLG 0357-3855	ESO 302- G 025	03 57 39.8	-38 55 53	14.95	1.47		S...	17618	
74	VLG 0405-0857		04 05 43.3	-08 57 53	13.95		10.60	21	9928	ir
75	VLG 0417-3917	ESO 303-IG 011	04 17 55.0	-39 17 30	14.93		9.96	Irr	15255	ir
76	VLG 0424-3440	ESO 360- G 009	04 24 46.2	-34 40 34	15.18	2.00		SA(rs)0-?	16031	
77	VLG 0430-3030	ESO 421- G 009	04 30 19.9	-30 30 15	14.98	1.40	10.56	S(r)a Lin	16481	ir,an
78	VLG 0435-3944	ESO 304- G 006	04 35 07.5	-39 44 45	14.71	0.91		SAB(rs)b	14283	
79	VLG 2105-3846	ESO 342- G 007	21 05 26.4	-38 46 06	15.30	0.41		Sa-b	17162	
80	VLG 2109-3919	ESO 342- G 020	21 09 25.3	-39 19 56	15.26	1.34		Sc	17159	
81	VLG 2109-3804	ESO 342- G 022	21 09 47.4	-38 04 33	14.50	0.92	10.48	Sb	12944	ir
82	VLG 2126-2124	ESO 599- G 016	21 26 47.6	-21 24 54	14.04	2.15		SA0-:	10608	rs
83	VLG 2157-1926		21 57 19.8	-19 26 41	15.48			E	17473	
84	VLG 2158-1507		21 58 02.2	-15 07 27	14.92			5	17096	
85	VLG 2205-1919	ESO 601- G 021	22 05 37.5	-19 19 06	14.67	1.43	9.87	SB(r)bc	14448	ir
86	VLG 2206-2803	ESO 467- G 012	22 06 17.2	-28 03 20	13.05	1.08	10.11	SB(s)bc p:Sy	6832	ir,an
87	VLG 2210-1653		22 10 22.9	-16 53 21	14.72		10.11	3	13403	ir
88	VLG 2214-3711		22 14 10.2	-37 11 52	15.08			5	17471	
89	VLG 2214-0455		22 14 20.5	-04 55 08	15.42		10.44	-5	17259	ir
90	VLG 2217-2033		22 17 18.9	-20 33 59	14.79		10.38	5	15338	ir

TABLE 1. (continued)

No.	Catalogue Name	Other Name	RA	Dec	m_B	B-R	$\log_{10}L_{60}$	Type (km/s)	$V_\odot$	Notes
91	VLG 2217-3256	ESO 405- G 009	22 17 33.4	-32 56 17	15.10	1.08		Sa	16992	
92	VLG 2222-1831		22 22 37.1	-18 31 24	15.05		10.17	5	15188	ir
93	VLG 2227-3046	ESO 468- G 003	22 27 01.4	-30 46 51	14.73	0.97		Sb	16021	
94	VLG 2229-2539	ESO 533- G 044	22 29 22.3	-25 39 20	13.92	1.47		SAB(rs)p	10157	
95	VLG 2231-3939		22 31 00.2	-39 39 22	14.98		9.98	S	17150	ir
96	VLG 2253-3449	ESO 406- G 022	22 53 05.7	-34 49 19	14.85	0.46		N	15519	
97	VLG 2256-2547	ESO 535- G 001	22 56 19.0	-25 47 47	13.84		9.85	(R:)SA(rs)a	9091	ir
98	VLG 2323-3809		23 23 52.6	-38 09 29	14.92			S	16135	
99	VLG 2326-3811		23 26 51.3	-38 11 18	15.06			S	15933	
100	VLG 2327-3924	ESO 347- G 019	23 27 34.6	-39 24 13	15.21	1.43		SB(r)ab	16611	
101	VLG 2327-3933		23 27 48.9	-39 33 26	14.79			E	16451	rs
102	VLG 2329-3933		23 29 07.9	-39 33 39	15.05			-2	15644	
103	VLG 2329-3419	ESO 408- G 008	23 29 11.2	-34 19 54	15.06	1.77		SAB(rs)	16234	
104	VLG 2329-3949		23 29 22.9	-39 49 52	15.13			S	16413	
105	VLG 2331-3837		23 31 46.2	-38 37 15	15.16			(R')SB(r)b	16255	
106	VLG 2332-3823	ESO 347- G 024	23 32 32.3	-38 23 54	15.08	1.43		S0-a	16387	
107	VLG 2338-2935		23 38 52.8	-29 35 53	15.05			E	15608	rs
108	VLG 2345-2825	ESO 471- G 019	23 45 08.3	-28 25 03	13.78?	1.43		cD;E+4 pec	8695	
109	VLG 2347-2917		23 47 38.1	-29 17 13	14.94			E	14580	
110	VLG 2348-3443	ESO 349- G 001	23 48 54.5	-34 43 50	14.52	1.81		E+1	17239	
111	VLG 2352-3452	ESO 349- G 005	23 52 22.6	-34 52 47	14.76	1.19		Sc	16002	
112	VLG 2354-3502	ESO 349- G 010	23 54 25.9	-35 02 16	14.29	1.70		cD;E+4;Lin	14696	rs,an
113	VLG 2356-2104	ESO 538- G 014	23 56 34.3	-21 04 35	15.02	1.14		Sa:	15140	

arXiv:astro-ph/9803033v1 4 Mar 1998

TABLE 2.

No .	Catalogue name	System type	System name	σ (km/s)	N_g
1	VLG 0006-3408	group candidate	SSRS2[a]		
4	VLG 0036-1541	triplet; interacting	IC 0038		
6	VLG 0039-0934	rich cluster; cD	A0085		
7	VLG 0041-2522	binary	ESO 474- G 020		
14	VLG 0049-2735	triplet	AM 0049-273		
17	VLG 0053-1015	rich cluster, binary	A0117		
18	VLG 0053-3740	binary	S0102		
20	VLG 0054-1011	group candidate	SSRS2[a]	907	9
21	VLG 0056-1644	binary	NGC0333A/B		
22	VLG 0056-2834	group, binary	SCG55		
23	VLG 0057-1415	rich cluster	A0126		
26	VLG 0100-2209	rich cluster; cD pec	A0133		
30	VLG 0106-1606	group	SSRS2[b]		
31	VLG 0106-1540	rich cluster	A0151		
32	VLG 0111-3200	poor cluster	S0141	292	17
35	VLG 0125-0456	compact group	HCG12		
36	VLG 0130-1504	binary; interacting	IC0141		
37	VLG 0131-3644	group	SSRS2[b]	68	5
38	VLG 0132-3644	group	SSRS2[b]	150	7
46	VLG 0148-0418	binary; interacting	NGC0702		
47	VLG 0150-1901	triplet	ESO 543- G 022		
48	VLG 0152-2823	binary	ESO 414-IG 009		
49	VLG 0154-2017	group	SCG14		
51	VLG 0159-2200	group	VV583		
52	VLG 0210-2242	group	SSRS2[b]		
54	VLG 0212-2504	rich cluster, triplet	A2995		
56	VLG 0215-2336	group	SSRS2[b]	877	9
57	VLG 0222-1922	group Seyfert	SSRS2[b]	146	6
59	VLG 0224-2338	poor cluster; brightest member	S0263		
60	VLG 0227-1715	poor cluster, binary	S0268		
62	VLG 0229-2006	group	SSRS2[b]		
65	VLG 0253-1424	group candidate; extended X source	SSRS2[a]		
67	VLG 0318-3842	binary	ESO 301-IG 006		
68	VLG 0322-0621	group	SSRS2[b]		

TABLE 2. (continued)

No .	Catalogue name	System type	System name	σ (km/s)	N_g
69	VLG 0327-2856	binary; interacting	ESO 418- G 007		
70	VLG 0337-3341	binary; interacting	ESO 358-IG 047		
71	VLG 0340-3811	group; interacting	AM 0340-381		
74	VLG 0405-0857	binary	IRAS 04057-0857		
75	VLG 0417-3917	group; interacting	ESO 303-IG 011		
80	VLG 2109-3919	binary	ESO 342- G 020		
86	VLG 2206-2803	compact group; Seyfert	HCG91 (VV700)	180	11
88	VLG 2214-3711	group	AM 2214-371		
91	VLG 2217-3256	triplet	AM 2217-325		
94	VLG 2229-2539	poor cluster	EDCC155	57	3
95	VLG 2231-3939	poor cluster	S1036		
97	VLG 2256-2547	group candidate	SSRS2[a]		
100	VLG 2327-3924	rich cluster	A4008	430	4
101	VLG 2327-3933	rich cluster; dumbbell system	A4008		
102	VLG 2329-3933	rich cluster	A4008		
103	VLG 2329-3419	poor cluster	EDCC312		
104	VLG 2329-3949	rich cluster	A4008		
107	VLG 2338-2935	poor cluster	EDCC332		
108	VLG 2345-2825	rich cluster; cD pec	A4038	307	16
112	VLG 2354-3502	rich cluster; cD , Liner	A4059		

[a]SSRS2 group candidate found in this work.
[b]SSRS2 group listed in the Girardi et al. catalog (1998).